

## Secondary Smoke Exposure At Home (School-age)

### Secondary Smoke Exposure At Home (School-age)

- Most recent data (2014): 20% of King County youth attending public schools in the 8th,10th and 12th grades.
- Trend over time: 2004-14: falling
- Description: Anyone who lives with you now smokes cigarettes
- Healthy People 2020 Goal: 41 percent or lower

Secondary smoke exposure at home (school-age), King County, 2012 and 2014 average

	Percent	Lower CI	Upper CI
<b>King County</b> (school-age in 8th, 10th and 12th grades)	21	21	22
<b>GRADE</b>			
Grade 8	20	19	22
Grade 10	22	20	24
Grade 12	22	20	24
<b>GENDER</b>			
Male	21	20	22
Female	22	21	23
<b>RACE/ETHNICITY</b>			
AIAN	36	31	41
Asian	22	21	24
Black	26	24	29
Hispanic	20	19	22
Multiple	25	23	27
NHPI	41	36	46
White	19	18	20
Other	23	21	26
<b>REGION</b>			
East	16	14	17
North	17	16	19
Seattle	22	20	25
South	27	26	28
<b>STATE</b>			
Washington State	26	25	27

#### Comparisons:

Higher than King County rate (yellow)

Lower than King County rate (green)

#### Notes:

Source: Healthy Youth Survey.

Prepared by Public Health - Seattle & King County, APDE, 08/2016.


CI is 95% Confidence Interval.

\* Too few cases to protect confidentiality and/or report reliable rates.

§ Too few cases to meet precision standard, interpret with caution.

Data by income or poverty level not available.

## Secondary smoke exposure at home (school-age) King County, 2012 and 2014 average



Source: Healthy Youth Survey.

Prepared by Public Health - Seattle & King County, APDE, 08/2016.


[-----] Confidence interval shows range that includes true value 95% of the time.

\* Too few cases to protect confidentiality and/or report reliable rates.

§ Too few cases to meet precision standard, interpret with caution.


Data by income or poverty level not available.

**Secondary smoke exposure at home (school-age) by regions  
King County, 2012 and 2014 average**



Source: Healthy Youth Survey.  
Prepared by Public Health - Seattle & King County, APDE, 08/2016.  
N/A: Not available.

## Secondary smoke exposure at home (school-age) King County, 2004-2014



Trends over time: King County (2004-14: falling); East KC (2004-14: falling); North KC (2004-14: falling); Seattle (2004-14: falling);  
 South KC (2004-14: falling); Washington State (2004-14: falling)  
 Prepared by Public Health - Seattle & King County, APDE, 08/2016.  
 Source: Healthy Youth Survey.  
 South KC (2004-14: falling); Washington State (2004-14: falling)

Produced by the  
[Assessment, Policy Development & Evaluation Unit](#)  
[Public Health - Seattle & King County](#)  
 08/16